

# Assessment of functional capacity before major non-cardiac surgery: an international, prospective cohort study



Duminda N Wijeyesundera, Rupert M Pearse, Mark A Shulman, Tom E F Abbott, Elizabeth Torres, Althea Ambosta, Bernard L Croal, John T Granton, Kevin E Thorpe, Michael P W Grocott, Catherine Farrington, Paul S Myles, Brian H Cuthbertson, on behalf of the METS study investigators

## Summary

**Background** Functional capacity is an important component of risk assessment for major surgery. Doctors' clinical subjective assessment of patients' functional capacity has uncertain accuracy. We did a study to compare preoperative subjective assessment with alternative markers of fitness (cardiopulmonary exercise testing [CPET], scores on the Duke Activity Status Index [DASI] questionnaire, and serum N-terminal pro-B-type natriuretic peptide [NT pro-BNP] concentrations) for predicting death or complications after major elective non-cardiac surgery.

**Methods** We did a multicentre, international, prospective cohort study at 25 hospitals: five in Canada, seven in the UK, ten in Australia, and three in New Zealand. We recruited adults aged at least 40 years who were scheduled for major non-cardiac surgery and deemed to have one or more risk factors for cardiac complications (eg, a history of heart failure, stroke, or diabetes) or coronary artery disease. Functional capacity was subjectively assessed in units of metabolic equivalents of tasks by the responsible anaesthesiologists in the preoperative assessment clinic, graded as poor (<4), moderate (4–10), or good (>10). All participants also completed the DASI questionnaire, underwent CPET to measure peak oxygen consumption, and had blood tests for measurement of NT pro-BNP concentrations. After surgery, patients had daily electrocardiograms and blood tests to measure troponin and creatinine concentrations until the third postoperative day or hospital discharge. The primary outcome was death or myocardial infarction within 30 days after surgery, assessed in all participants who underwent both CPET and surgery. Prognostic accuracy was assessed using logistic regression, receiver-operating-characteristic curves, and net risk reclassification.

**Findings** Between March 1, 2013, and March 25, 2016, we included 1401 patients in the study. 28 (2%) of 1401 patients died or had a myocardial infarction within 30 days of surgery. Subjective assessment had 19·2% sensitivity (95% CI 14·2–25) and 94·7% specificity (93·2–95·9) for identifying the inability to attain four metabolic equivalents during CPET. Only DASI scores were associated with predicting the primary outcome (adjusted odds ratio 0·96, 95% CI 0·83–0·99;  $p=0\cdot03$ ).

**Interpretation** Subjectively assessed functional capacity should not be used for preoperative risk evaluation. Clinicians could instead consider a measure such as DASI for cardiac risk assessment.

**Funding** Canadian Institutes of Health Research, Heart and Stroke Foundation of Canada, Ontario Ministry of Health and Long-Term Care, Ontario Ministry of Research, Innovation and Science, UK National Institute of Academic Anaesthesia, UK Clinical Research Collaboration, Australian and New Zealand College of Anaesthetists, and Monash University.

Copyright © 2018 Elsevier Ltd. All rights reserved.

## Introduction

International clinical practice guidelines emphasise the assessment of preoperative cardiopulmonary fitness, or functional capacity, as an important component of estimating patients' risks for major morbidity and mortality after surgery.<sup>1,2</sup> For example, the American College of Cardiology and American Heart Association recommend that patients proceed directly to elective intermediate and major non-cardiac surgery if they are capable of more than four metabolic equivalents of tasks of activity without symptoms.<sup>1</sup> The usual standard of care for assessing preoperative functional capacity involves doctors assessing patients, then making subjective estimates of their fitness (ie, subjective assessment). Although these estimates are easily implementable into clinical practice,

subjective assessment has limitations, including little agreement with validated measures of functional capacity,<sup>3</sup> and poor accuracy when used to predict post-operative death or complications.<sup>4,5</sup> These limitations point to the need for better alternatives to assess preoperative functional capacity.

Possible alternative options are cardiopulmonary exercise testing (CPET), which has been described as a gold standard non-invasive assessment of exercise tolerance, and the Duke Activity Status Index (DASI),<sup>6</sup> which is a standardised questionnaire correlated with gold-standard measures of functional capacity. Additionally, although no blood test can directly measure functional capacity, N-terminal pro-B-type natriuretic peptide (NT pro-BNP) concentrations might indirectly fulfil this role.<sup>7</sup>

*Lancet* 2018; 391: 2631–40

See [Comment](#) page 2580

Li Ka Shing Knowledge Institute (D N Wijeyesundera PhD, K E Thorpe MMath) and Applied Health Research Centre (E Torres BSc, A Ambosta MA, K E Thorpe), St Michael's Hospital, Toronto, ON, Canada; Department of Anesthesia and Pain Management (D N Wijeyesundera) and Department of Medicine (Prof J T Granton MD), University Health Network, Toronto, ON, Canada; Department of Anesthesia (D N Wijeyesundera, Prof B H Cuthbertson MD), Institute of Health Policy Management and Evaluation (D N Wijeyesundera, Prof B H Cuthbertson), Department of Medicine (Prof J T Granton), and Dalla Lana School of Public Health (K E Thorpe), University of Toronto, Toronto, ON, Canada; Department of Medicine, Sinai Health System, Toronto, ON, Canada (Prof J T Granton); Department of Critical Care Medicine, Sunnybrook Health Sciences Centre, Toronto, ON, Canada (Prof B H Cuthbertson); Queen Mary University of London, London, UK (Prof R M Pearse MD, T E F Abbott PhD); NHS Grampian, Aberdeen, UK (B L Croal MD); University Hospital Southampton, Southampton, UK (Prof M P W Grocott MD), University of Southampton, Southampton, UK (Prof M P W Grocott); Alfred Hospital, Melbourne, VIC, Australia (M A Shulman MPH, C Farrington BAppSc, Prof P S Myles DSc); and Monash University, Melbourne, VIC, Australia (M A Shulman, C Farrington, Prof P S Myles)

Correspondence to:  
Prof Brian H Cuthbertson,  
Department of Critical Care  
Medicine, Sunnybrook Health  
Sciences Centre, Toronto,  
ON M4N 3M5, Canada  
brian.cuthbertson@  
sunnybrook.ca

### Research in context

#### Evidence before this study

Estimation of cardiopulmonary fitness, or functional capacity, is an important component of risk assessment before major non-cardiac surgery. This estimation typically involves subjective assessment, where doctors interview patients and make a subjective judgment of their fitness. To assess the validity of this commonly used measure of functional capacity, we used the terms [("prediction" OR "preoperative evaluation" OR "risk prediction") AND ("surgery" AND "complications") AND ("exercise capacity" OR "activities of daily living" OR "functional capacity")] to search the PubMed database for English-language articles on relevant studies published before Dec 31, 2017. The search was supplemented with hand-searches of reference lists from relevant reviews and practice guidelines. Previous research was limited to single-centre studies with small sample sizes or a high risk of bias. In these studies, subjective assessment showed poor agreement with validated questionnaires, and an inconsistent association with postoperative complications.

#### Added value of this study

Our multicentre prospective cohort study (Measurement of Exercise Tolerance before Surgery) assessed patients before they underwent major elective non-cardiac surgery and

compared the prognostic accuracy of subjective assessment against three alternative measures: the Duke Activity Status Index (DASI) questionnaire, cardiopulmonary exercise testing (CPET) to measure peak oxygen consumption, and serum N-terminal pro-B-type natriuretic peptide (NT pro-BNP) concentrations. In a sample of 1401 adult participants at 25 hospitals, lower DASI scores predicted 30-day death or myocardial infarction, and 30-day death or myocardial injury; higher NT pro-BNP concentrations predicted 30-day death or myocardial injury, and 1-year death; and lower peak oxygen consumption predicted complications. Subjective assessment did not predict any outcomes.

#### Implications of all the available evidence

Subjective assessment of functional capacity should not be used for preoperative risk assessment. This commonly used practice does not accurately identify patients with poor fitness or those at increased risk for postoperative morbidity and mortality. As alternatives, clinicians could consider more objective measures, such as DASI questionnaires and NT pro-BNP testing to assess perioperative cardiac risk, and perhaps CPET to predict complications after major elective non-cardiac surgery.

Increased NT pro-BNP concentrations are integrated markers of cardiac dysfunction, including myocardial stretch and ischaemia. We therefore did a study to compare preoperative subjective assessment, CPET, DASI, and NT pro-BNP for predicting death or complications after major elective non-cardiac surgery.

## Methods

### Study design and participants

We did a multicentre, international, prospective cohort study (Measurement of Exercise Tolerance before Surgery [METS]) at 25 hospitals: five in Canada, seven in the UK, ten in Australia, and three in New Zealand. The study's objectives, design, and methods have been previously reported.<sup>8</sup> Details of the methods are in the appendix.

Participants were aged 40 years or older, scheduled for elective non-cardiac surgery under general or regional anaesthesia (or both) with a minimum of one overnight hospital stay, and deemed to have at least one risk factor for cardiac complications or at least one risk factor for having coronary artery disease (appendix). All participants provided written informed consent, and each centre obtained research ethics board approval before commencing recruitment.

### Procedures

During the period from recruitment to 1 day before surgery, participants underwent symptom-limited incremental CPET on a computer-controlled, electro-magnetically braked cycle ergometer using a standardised

protocol (appendix).<sup>8</sup> This assessment usually occurred during a separate hospital visit after the date of recruitment. On the basis of a prespecified assessment of the plotted CPET data, trained investigators at each centre determined patients' peak oxygen consumption and anaerobic threshold.

Participants also underwent three other preoperative assessments of functional capacity. First, responsible anaesthesiologists in the preoperative assessment clinic (on the date of recruitment) or operating theatre (on the day of surgery) were asked to make a subjective judgment of participants' functional capacity after assessing their usual preoperative history. Subjectively assessed functional capacity was classified as poor (<4 metabolic equivalents), moderate (4–10 metabolic equivalents), or good (>10 metabolic equivalents). The poor category included cases where anaesthesiologists were uncertain about patients' functional capacity, typically due to pre-existing conditions such as arthritis or peripheral arterial disease.<sup>1</sup> Second, participants completed the DASI questionnaire on the date of recruitment (appendix). Third, blood samples were drawn at any point between recruitment and surgery to measure serum NT pro-BNP concentrations. These samples were stored at –70°C to –80°C at each study site, then analysed at the Aberdeen Royal Infirmary (Aberdeen, UK) using the Siemens Vista immunoassay analyser (Siemens Healthcare Diagnostics, Frimley, UK). Participants, health-care providers, and outcome adjudicators were masked to CPET and NT pro-BNP results, and health-care providers and outcome adjudicators were masked to DASI scores; specifically,

See Online for appendix

the study's central laboratory, exercise testing facilities, and exercise testing laboratories did not report these specific measurements to masked individuals. The exceptions were cases of myocardial ischaemia or substantial new arrhythmias during CPET, in which case these specific findings, but not peak oxygen consumption or anaerobic threshold results, were revealed to health-care providers.

After surgery, participants underwent daily electrocardiograms and blood sampling to measure troponin and creatinine concentrations, until the third post-operative day or hospital discharge (whichever came first). Research personnel followed participants each day throughout their hospital stay to ascertain the presence of specific complications (appendix). The severity of complications was categorised as mild, moderate, severe, or fatal using a modified Clavien-Dindo classification scheme.<sup>9,10</sup> After hospital discharge, participants were contacted at 30 days and 1 year after surgery to ascertain vital status. Details of the follow-up process are in the appendix.

### Outcomes


The primary outcome was death or myocardial infarction within 30 days after surgery. The secondary outcome was death within 1 year after surgery. Myocardial infarction was diagnosed by an adjudication committee that used the third universal definition of myocardial infarction<sup>11</sup> while remaining masked to CPET, DASI, and NT pro-BNP results. Other outcomes of interest were death or myocardial injury within 30 days after surgery, and moderate or severe (including fatal) complications during the index admission to hospital. Myocardial injury was defined as postoperative troponin concentrations exceeding both the 99th percentile of the normal reference population, and the threshold at which the assay coefficient of variation was 10%. Moderate or severe complications were included as an endpoint because these events have been associated with poor preoperative functional capacity, especially when measured objectively by CPET.<sup>12,13</sup>

### Statistical analysis

The sample size calculation was based on comparing the area under the curve (AUC) of the receiver-operating-characteristic (ROC) curves for peak oxygen consumption versus subjective assessment with respect to predicting myocardial infarction or death within 30 days of surgery. During the design of the study, we initially calculated that a sample size of 1180 was required on the basis of underlying assumptions of an outcome event rate of 8%, correlation of 0.5 between peak oxygen consumption and subjective assessment,<sup>6</sup> an AUC of 0.65 for subjective assessment, an AUC of 0.75 for peak oxygen consumption, and 90% power to detect this difference in AUC values (two-sided  $\alpha$  of 0.05). To account for 10% of participants not undertaking CPET or surgery, we aimed to recruit 1312 participants. After recruiting about half

the original planned sample size, this calculation was reassessed based on two factors identified in the accumulating study data. First, we noted that about 20% of participants did not either undertake CPET or undergo their planned surgeries. Second, the event rate for the primary outcome was instead projected to be approximately 5%. The overall sample size was therefore increased to 1723 participants to account for up to 20% of recruited individuals not being eligible for the primary analysis, and a primary outcome event rate of 5%, while retaining the power of 80%. We remained masked to all data on the principal exposures (ie, CPET results, DASI scores, and NT pro-BNP concentrations) during the sample size re-estimation.

All participants who undertook both CPET and surgery were included in the primary analysis; CPET performance was characterised by peak oxygen consumption. For each outcome of interest, we built separate nested logistic regression models that sequentially included baseline clinical characteristics followed by the exposure of interest



**Figure 1: Participant screening, recruitment, and follow-up**  
CPET=cardiopulmonary exercise testing.

	Full cohort (n=1401)	Missing data
<b>Demographic characteristics</b>		
Age (years)	65 (57–72)	..
Female sex	548 (39%)	..
<b>Comorbidities</b>		
Coronary artery disease	165 (12%)	..
Heart failure	20 (1%)	..
Cerebrovascular disease	59 (4%)	..
Peripheral arterial disease	42 (3%)	..
Diabetes mellitus	264 (19%)	..
Hypertension	779 (56%)	..
Current or recent smoker*	216 (15%)	..
Obstructive lung disease†	181 (13%)	..
Significant arthritis‡	289 (21%)	..
Significant malignancy§	597 (43%)	..
Preoperative renal function¶		49
eGFR ≥60 mL/min per 1.73 m <sup>2</sup>	1195 (88%)	..
eGFR 30–59 mL/min per 1.73 m <sup>2</sup>	125 (9%)	..
eGFR <30 mL/min per 1.73 m <sup>2</sup> or dialysis	31 (2%)	..
<b>Composite risk scales</b>		
ASA-PS classification		3
Class 1	103 (7%)	..
Class 2	818 (59%)	..
Class 3	457 (33%)	..
Class 4	20 (1%)	..
Revised Cardiac Risk Index		..
Class 1	624 (45%)	..
Class 2	635 (45%)	..
Class 3	115 (8%)	..
Class 4	27 (2%)	..
<b>Preoperative medications</b>		
β blocker	232 (17%)	..
Dihydropyridine calcium channel blocker	248 (18%)	..
Diltiazem or verapamil	27 (2%)	..
ACE inhibitor or ARB	529 (38%)	..
Furosemide	55 (4%)	..
Aspirin	334 (24%)	..
Other antiplatelet medication	35 (3%)	..

(Table 1 continues in next column)

(ie, subjective assessment, peak oxygen consumption, anaerobic threshold, DASI scores, or NT pro-BNP). We modelled NT pro-BNP concentrations using a logarithmic transformation to reduce the potential effect of extreme values within its highly skewed distribution. The statistical significance of prognostic information from additional predictors was based on the increase in log likelihood of the larger model. For the models predicting the primary and secondary outcomes, the baseline variable was the validated Revised Cardiac Risk Index (RCRI) score.<sup>14,15</sup> In the model predicting death or myocardial injury by 30 days, the baseline variables were age, sex, and RCRI score. The baseline variables in the model predicting moderate-or-severe complications were

	Full cohort (n=1401)	Missing data
(Continued from previous column)		
<b>Operative characteristics</b>		
Procedure type		
Vascular	26 (2%)	..
Intrathoracic	31 (2%)	..
Intraperitoneal or retroperitoneal	464 (33%)	..
Urological or gynaecological	417 (30%)	..
Head and neck	93 (7%)	..
Orthopaedic	336 (24%)	..
Other	30 (2%)	..
Laparoscopic or thoracoscopic assistance	499 (36%)	..
Anaesthesia type		
General anaesthesia alone	758 (54%)	..
Regional anaesthesia alone	210 (15%)	..
General plus regional anaesthesia	433 (31%)	..
Intraoperative haemodynamic monitoring		
Arterial line	687 (49%)	7
Central venous line	212 (15%)	7
Cardiac output monitor	95 (7%)	8
Postoperative disposition		
Critical-care unit or monitored bed unit	331 (24%)	1

Data are median (IQR), n (%), or ..=not applicable. eGFR=estimated glomerular filtration rate. ASA-PS=American Society of Anesthesiologists Physical Status. ACE=angiotensin-converting enzyme. ARB=angiotensin-receptor blocker. \*Current smoker or quit within previous 1 year. †Previous diagnosis of asthma, reactive airways disease, chronic obstructive lung disease, chronic bronchitis, or emphysema. ‡Previous or scheduled major joint replacement surgery. §Indication for surgery was for treatment of cancer. ¶eGFR was calculated using the preoperative serum creatinine concentration and Chronic Kidney Disease Epidemiology Collaboration equation. ||Revised Cardiac Risk Index scores were calculated using the original definitions of diabetes (ie, requirement for insulin therapy) and renal insufficiency (ie, creatinine concentration >176 μmol/L).<sup>15</sup> When determining Revised Cardiac Risk Index scores, any individual with missing preoperative creatinine concentration data was assumed to have a concentration of 176 μmol/L or less.

**Table 1: Baseline characteristics of study cohort**

age, sex, and high-risk surgery, which was defined as intraperitoneal, intrathoracic, or suprainguinal vascular procedures.<sup>14</sup> These covariates were selected a priori based on previous evidence, their inclusion in guideline-recommended assessment algorithms,<sup>1,16</sup> and the need to prevent model overfitting. Additionally, the covariates mirror clinically sensible factors typically considered during preoperative assessment. We calculated the area under the ROC curve of models with successively more predictors, and models with only the individual exposure of interest (eg, peak oxygen consumption). Prognostic information from these models was compared using the continuous net reclassification improvement (NRI) statistic and area under the ROC curve.<sup>17</sup>

To address whether preoperative functional capacity might have better prognostic value in more invasive surgical procedures, a post-hoc subset analysis was done in some patients undergoing body cavity surgery, which was defined as intrathoracic, intraperitoneal,

retroperitoneal or pelvic (ie, urological or gynaecological) procedures.

Analyses were done with R (version 3.4.0), statistical significance was defined by a two-tailed p value of less than 0.05, and no adjustments were made for multiple comparisons.<sup>18</sup> Because missing data were uncommon, a complete case analysis was done. Additionally, these missing data pertained to baseline information that was likely missing completely at random.

### Role of the funding source

The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. The corresponding author had full access to all the data in the study and had final responsibility for the decision to submit for publication.

### Results

Between March 1, 2013, and March 25, 2016, 1741 patients (23% of eligible patients who consented to participate) were recruited at 25 hospitals, with 1401 (81%) undergoing CPET and surgery (figure 1). Of these 1401 participants in the primary cohort, 1399 (99.9%) completed a 30-day follow-up and 1378 (98%) completed a 1-year follow-up. Participants' median age was 65 years (IQR 57–72), 39% were female, and 91% were classified as American Society of Anesthesiologists Physical Status (ASA-PS) 2 or 3 (table 1). Most participants underwent major abdominal, pelvic, or orthopaedic procedures. Data were missing for less than 4% of participants.

The CPET exercise protocol was terminated early in 11% of participants, with the most common reasons being inability to pedal, fatigue, or a safety-based indication (table 2). Mean peak oxygen consumption was 19.2 mL/kg per min (SD 6.5) and mean anaerobic threshold was 12.6 mL/kg per min (4.1). 110 (8%) of participants had adverse events during CPET (table 2), with 27 cases meeting criteria for unmasking. The clinical sequelae of these events are in the appendix.

Additionally, 1351 (96%) participants underwent subjective assessment of functional capacity, 1396 (99.6%) completed DAS1 questionnaires, and 1347 (96%) had NT pro-BNP concentrations measured (appendix). 107 (8%) participants were subjectively judged to have poor preoperative functional capacity, 230 (16%) had a peak oxygen consumption value of less than 14 mL/kg per min (equivalent to <4 metabolic equivalents), and 426 (30%) had an anaerobic threshold below the suggested high-risk threshold of 11 mL/kg per min.<sup>19</sup> The characteristics of participants within strata defined by peak oxygen consumption values is in the appendix. A subjective assessment of poor functional capacity had a sensitivity of 19.2% (95% CI 14.2–25.0) for identifying peak oxygen consumption of less than 14 mL/kg per min, while its specificity was 94.7% (93.2–95.9).

When stratified by subjectively assessed functional capacity, peak oxygen consumption and DAS1 values were

	Overall cohort (n=1401)
Interval to CPET to surgery (days)	9 (5–21)
Early termination of CPET	157 (11%)
Reasons for early termination of CPET	
Safety-based indication	23 (2%)
Fatigue	31 (2%)
Shortness of breath	11 (<1%)
Unable to pedal	76 (5%)
Unable to tolerate mouthpiece or mask	12 (<1%)
Technical problems with equipment	4 (<1%)
Measurable peak oxygen consumption	1356 (97%)
Measurable AT	1275 (91%)
Adverse events during CPET	
Any adverse event	110 (8%)
Ischaemic ECG changes	25 (2%)
Chest pain	2 (<1%)
Significant new arrhythmias	14 (1%)
Significant hypertension	43 (3%)
Significant drop in blood pressure	28 (2%)
Syncope	3 (<1%)
Significant decrease in oxygen saturation	10 (<1%)

Data are median (IQR) or n (%). CPET=cardiopulmonary exercise testing. AT=anaerobic threshold. ECG=electrocardiogram.


**Table 2: Characteristics of CPET assessments**

generally lower in individuals judged to have poor functional capacity, but there was substantial within-stratum variation (figure 2). Peak oxygen consumption was positively correlated with DAS1 scores (figure 2; Spearman  $\rho$  0.43,  $p < 0.0001$ ), and negatively correlated with NT pro-BNP concentrations (Spearman  $\rho$  -0.21,  $p < 0.0001$ ). There was also a negative correlation between DAS1 scores and NT pro-BNP concentrations (Spearman  $\rho$  -0.25,  $p < 0.0001$ ; appendix).

After surgery, 194 (14%) participants had in-hospital moderate-or-severe complications.

By 30 days after surgery, five (<1%) participants had died, 24 (2%) had a myocardial infarction, 28 (2%) had the primary outcome of death or myocardial infarction, and 176 (13%) had died or had a myocardial injury. By 1 year after surgery, 38 participants (3%) had died. Of the moderate or severe complications, the more frequent events were respiratory failure, pneumonia, surgical site infection, re-operation, and unexpected critical care unit admission (appendix).

Subjectively assessed preoperative functional capacity had no significant adjusted association with the four main study outcomes (appendix). A significant adjusted association and significant risk reclassification with peak oxygen consumption was observed only with respect to moderate or severe complications (table 3). Anaerobic threshold showed no significant association or risk reclassification with the main outcomes. DAS1 scores showed significant adjusted associations with the primary outcome of death or myocardial infarction by



**Figure 2: Correlation between measures of preoperative functional capacity**

(A) is a boxplot presenting the distributions of peak oxygen consumption within strata defined by subjectively assessed preoperative functional capacity, categorised as poor (<4 metabolic equivalents), moderate (4–10 metabolic equivalents) or good (>10 metabolic equivalents). The horizontal line within each box denotes the median, and the top and bottom of each box indicate the IQR. Vertical lines at the top and bottom of each box extend to the 5th and 95th percentiles, and solid circles indicate outliers. 4 metabolic equivalents correspond to a peak oxygen consumption of 14 mL/kg per min, and 10 metabolic equivalents correspond to a peak oxygen consumption of 35 mL/kg per min. The boxplot on the right (denoted as not applicable [NA]) presents the peak oxygen consumption distribution among the 50 participants with missing subjective assessments. B is a boxplot presenting the distributions of scores on the Duke Activity Status Index (DASI) within strata defined by subjectively assessed preoperative functional capacity. The minimum possible DASI score is 0 and the maximum possible score is 58.2. The boxplots on the right (denoted as NA) presents the DASI score distribution among the 50 participants with missing subjective assessments. The wide-ranging distribution in panels (A) and (B) for participants with missing assessments suggests that these values were missing completely at random. (C) is a scatter plot presenting the association of peak oxygen consumption (y axis) and DASI scores (x axis). The Spearman correlation coefficient between DASI scores and peak oxygen consumption was 0.41 ( $p < 0.0001$ ). (D) is a scatter plot presenting the association of peak oxygen consumption (y axis) and N-terminal pro-B-type natriuretic peptide (NT pro-BNP) concentrations (x axis on a  $\log_{10}$  scale). The Spearman correlation coefficient between NT pro-BNP and peak oxygen consumption was  $-0.21$  ( $p < 0.0001$ ). In both panels C and D, the line of best fit is shown in blue (estimated using cubic regression splines), and the grey shaded zones represent its 95% CIs.

30 days after surgery and with death or myocardial injury by 30 days after surgery. Additionally, DASI scores showed significant risk reclassification with death or myocardial injury by 30 days. NT pro-BNP concentrations showed significant adjusted associations and significant risk reclassification with death by 1 year after surgery and death or myocardial injury by 30 days after surgery. When the main study analyses were repeated in

the post-hoc subset, the results remained qualitatively unchanged (appendix).

## Discussion

Preoperative subjective assessment neither accurately identified patients with poor cardiopulmonary fitness nor predicted postoperative morbidity and mortality. The DASI questionnaire improved prediction of 30-day myocardial infarction or death, and 30-day myocardial injury or death; and NT pro-BNP concentrations improved prediction of 30-day myocardial injury or death, and 1-year death. Formal assessment of cardiopulmonary fitness, based on peak oxygen consumption during CPET, improved prediction of moderate or severe postoperative complications.

In our study, subjective assessment of preoperative functional capacity consistently performed poorly. Although it had construct validity, in that peak oxygen consumption was generally lower in patients judged to be less fit, subjective assessment correctly identified only 16% of patients who achieved a peak less than 14 mL/kg per min, which is consistent with less than 4 metabolic equivalents. Further, subjective assessment did not predict postoperative myocardial infarction, myocardial injury, or myocardial complications, confirming findings from a single-centre retrospective cohort study that relied on an administrative database for outcome ascertainment.<sup>5</sup> Based on these findings, subjective assessment should not be used to assess patients' risks of major postoperative cardiac complications.

Notably, more objective assessment of cardiopulmonary fitness with CPET did not improve most aspects of preoperative risk assessment. Consistent with previous evidence,<sup>12</sup> peak oxygen consumption measured during CPET was predictive of postoperative complications; however, most of these events were pulmonary complications, surgical site infections, unexpected critical care unit admissions, and re-operations. By contrast, peak oxygen consumption and anaerobic threshold were not associated with postoperative myocardial infarction or myocardial injury, somewhat contradicting the emphasis of practice guidelines on functional capacity for preoperative cardiac risk evaluation.<sup>12</sup> These findings occurred within the context of our study addressing several important limitations in the current evidence base,<sup>20</sup> in that it masked CPET results (unlike most previous studies), and implemented standardised outcome surveillance in a large, generalisable, multicentre sample. There are several possible explanations for our results. First, the previous evidence supporting a link between fitness and perioperative cardiac risk had limitations; eg, many of the studies were done more than 30 years ago, and have limited generalisability to contemporary patients and surgeries.<sup>21,22</sup> Other studies had few outcome events or associations of only weak magnitudes.<sup>23,24</sup> Second, low peak oxygen consumption or anaerobic threshold might not be the ideal CPET-based indicator of the underlying causal

mechanisms for perioperative myocardial infarction. Other metrics, such as an exaggerated exercise-mediated heart rate response,<sup>25</sup> might be better indicators of perioperative cardiac risk.

The DASI questionnaire had construct validity as a measure of functional capacity in surgical patients, consistent with previous research,<sup>26</sup> and also improved prediction of postoperative myocardial infarction and myocardial injury. Our findings confirmed non-operative data indicating enhanced risk prediction using this questionnaire,<sup>27</sup> supported guideline suggestions for using objective scales to assess functional capacity,<sup>1</sup> and indicate opportunities for straightforward improvements in clinical practice. Specifically, the simple DASI questionnaire can be easily implemented into most perioperative practice settings, although further studies are needed to define optimal risk-specific thresholds in DASI scores, and develop reliable non-English versions of the questionnaire.<sup>28</sup> An important area of residual uncertainty is why DASI scores were associated with postoperative cardiac events, yet peak oxygen consumption was not. Given the only moderate correlation between DASI scores and peak oxygen consumption, a possible explanation is that DASI also measures somewhat different constructs, such as musculoskeletal strength, frailty, and self-imposed physical limitations.<sup>29</sup>

Confirming results from a previous individual patient data meta-analysis,<sup>30</sup> increased preoperative NT pro-BNP concentrations were associated with increased risks of postoperative 30-day death or myocardial injury in the METS study cohort; additionally, increased concentrations also predicted increased 1-year mortality. These findings support recommendations in recent practice guidelines to incorporate natriuretic-peptide testing into preoperative risk assessment strategies.<sup>16</sup> We noted only slight-to-fair correlation between NT pro-BNP concentrations and measures of exercise capacity (ie, peak oxygen consumption and DASI). This low correlation suggests that NT pro-BNP measures are a construct distinct from exercise capacity, and raises the possibility of enhancing preoperative assessment by combining measures of functional capacity and NT pro-BNP in future risk prediction models.

Our study had several limitations. First, despite increasing the original projected sample size, the primary and secondary outcomes occurred fewer times than anticipated. To some extent, the risks of death within 30 days of surgery (ie, 0.4%) and 30-day death or myocardial infarction (ie, 2.0%) in our study are representative of risks for contemporary elective major non-cardiac surgery in high-income countries, as suggested by several studies published after the design of our study. For example, in the ISOS prospective cohort study of 44814 adults having elective inpatient surgery across 27 high-income and middle-income countries, the risk of in-hospital 30-day death was 0.5%.<sup>10</sup> Similarly, in

	Adjusted odds ratio (95% CI)	AUC*	Net reclassification improvement index†		
			Events	Non-events	Overall
<b>30-day death or myocardial infarction</b>					
Baseline model‡		0.59			
Plus peak oxygen consumption	0.90 (0.71–1.16; p=0.45)§	0.62	0.04	-0.04	0.01 (p=0.98)
Plus AT	0.96 (0.66–1.41; p=0.84)§	0.59	-0.24	-0.12	-0.36 (p=0.10)
Plus DASI	0.91 (0.83–0.99; p=0.03)§	0.67	0.07	0.21	0.28 (p=0.14)
Plus NT PRO-BNP	1.88 (0.89–3.96; p=0.09)¶	0.65	0.11	0.14	0.25 (p=0.19)
<b>30-day death or myocardial injury</b>					
Baseline model		0.70			
Plus peak oxygen consumption	1.03 (0.92–1.14; p=0.62)§	0.70	-0.07	0.16	0.09 (p=0.26)
Plus AT	1.12 (0.96–1.31; p=0.16)§	0.71	0.17	-0.08	0.09 (p=0.27)
Plus DASI	0.96 (0.92–0.99; p=0.05)§	0.71	0.05	0.19	0.23 (p=0.004)
Plus NT PRO-BNP	1.78 (1.21–2.62; p=0.003)¶	0.71	0.07	0.13	0.20 (p=0.02)
<b>In-hospital moderate or severe complications</b>					
Baseline model**		0.72			
Plus peak oxygen consumption	0.86 (0.78–0.97; p=0.007)§	0.74	0.21	-0.004	0.21 (p=0.008)
Plus AT	0.87 (0.74–1.02; p=0.08)§	0.69	0.17	-0.08	0.09 (p=0.27)
Plus DASI	0.97 (0.93–1.01; p=0.16)§	0.72	-0.17	0.15	-0.02 (p=0.80)
Plus NT PRO-BNP	1.10 (0.77–1.57; p=0.61)¶	0.72	0.03	0.08	0.12 (p=0.12)
<b>1-year death</b>					
Baseline model‡		0.65			
Plus peak oxygen consumption	0.94 (0.77–1.15; p=0.56)§	0.66	0.20	-0.06	0.14 (p=0.39)
Plus AT	1.03 (0.76–1.40; p=0.56)§	0.64	0.13	0.15	0.28 (p=0.12)
Plus DASI	0.94 (0.87–1.02; p=0.13)§	0.69	0.00	0.16	0.16 (p=0.34)
Plus NT PRO-BNP	2.91 (1.54–5.49; p=0.001)¶	0.72	0.17	0.23	0.39 (p=0.02)

AUC=area under the curve. AT=anaerobic threshold. DASI=Duke Activity Status Index. NT PRO-BNP=N-terminal pro-B-type natriuretic peptide. \*AUC of the receiver operating characteristic curve for the relevant logistic regression model. †The weighed net proportion of individuals whose predicted probability of the outcome of interest improved with inclusion of a specific additional covariate in the regression model (eg, peak oxygen consumption). Improved predicted probability implies a higher predicted probability in individuals who had the outcome event of interest, and a lower predicted probability in individuals who did not. Negative statistic values indicate net worsening of predicted probabilities (eg, higher predicted probability in individuals who did not have the outcome of interest). This statistic has an associated p value. ‡Covariate in this baseline model was the Revised Cardiac Risk Index score. §Adjusted odds ratios (OR) expressed with respect to 1 metabolic equivalent increase in peak oxygen consumption (per 3.5 mL/kg per min), AT (per 3.5 mL/kg per min), or DASI scores (per 3.5 points). ¶Adjusted OR expressed with respect to 1 log<sub>10</sub> increase in NT pro-BNP concentrations ||Covariates in this baseline model were age, sex, and Revised Cardiac Risk Index score. \*\*Covariates in this baseline model were age, sex, and high-risk surgery (intraoperative, intrathoracic, or suprainguinal vascular procedures).

**Table 3: Predictive performance of different measures of preoperative functional capacity**

the VISION prospective cohort study of 15133 patients having inpatient non-cardiac surgery in eight high-income and middle-income countries, the risk of 30-day mortality after elective surgery was 1.2%. In this same cohort, the overall risk of myocardial infarction was about 3.3%.<sup>31</sup> Since 14% of the cohort had emergency surgery, which is associated with a three-times higher risk of myocardial infarction,<sup>32</sup> the risk of myocardial infarction after elective surgery in the VISION study was probably about 2.6%. Thus, the event rates in our study were consistent with other contemporary major non-cardiac surgery samples. Nonetheless, to help to address this reduced statistical power, we analysed the association

of the exposures of interest with two more frequent outcomes: myocardial injury and moderate or severe postoperative complications. Myocardial injury and postoperative complications are clinically and prognostically important outcomes.<sup>32–34</sup> Our general findings with respect to prediction of 30-day myocardial infarction or death were qualitatively unchanged in the complementary analysis pertaining to 30-day myocardial injury or death.

Second, despite significant efforts by research personnel and study investigators, the consent rate among otherwise eligible patients in our study was only 27%. Nonetheless, this consent rate is somewhat unsurprising when viewed from the perspective of the study setting (ie, anxious patients awaiting major surgery within a short timeframe) and procedures (ie, strenuous exercise testing solely for research purposes). The consent rate is also consistent with several large contemporary prospective studies in surgical patients that had arguably more straightforward study procedures. For example, consent rates were 30% in the POISE-2 trial of aspirin and clonidine in non-cardiac surgery,<sup>35</sup> and 36% in the ATACAS trial in cardiac surgery.<sup>36</sup>

Third, our primary analyses relied on peak oxygen consumption and anaerobic thresholds determined by trained investigators at each individual centre. It is possible that central adjudication of CPET results might have led to different determinations of these measures. Nonetheless, given that our study was designed to be pragmatic and generalisable, our main analyses better represent the prognostic accuracy of peak oxygen consumption or anaerobic thresholds in real world clinical practice. Fourth, the preoperative predictive models in this study had generally low-to-moderate discrimination, with AUC values of 0.74 or lower. This observation could be partly explained by the lower-than-expected outcome event rate, which limited the number of covariates included in regression models. Nonetheless, the discrimination of these models is similar to those from other studies, such as a prospective cohort study where the combination of RCRI score and preoperative coronary CT angiography had an AUC of 0.66 for predicting 30-day death or myocardial infarction.<sup>37</sup> Fifth, each hospital used its own preferred troponin assay to detect myocardial infarction or myocardial injury. This pragmatic approach is consistent with many multicentre perioperative studies.<sup>35,38</sup> Furthermore, variation in troponin assays does not affect the prognostic importance of myocardial infarction or myocardial injury,<sup>32,33</sup> and should not have biased the association between the study exposures and outcomes.

Combined with previous data, the results of our study suggest that DASI scores and natriuretic peptides, such as NT pro-BNP, should supplant subjective assessment for the estimation of perioperative cardiac risk for major non-cardiac surgery. More research is required to define optimum thresholds for these measures and determine

how they should be used in combination with other prognostically important information, including alternative preoperative biomarkers (eg, high-sensitivity troponin).<sup>39</sup> These other biomarkers might also help to partly address the limitations of NT pro-BNP as a prognostic biomarker in the presence of obesity,<sup>40</sup> chronic renal kidney,<sup>41</sup> or heart failure with preserved ejection fraction.<sup>42</sup>

Our findings also indicate that CPET, specifically peak oxygen consumption, can be used to identify patients at increased risk for postoperative complications. Research is needed to define optimal thresholds in peak oxygen consumption, determine the role of central expert adjudication in improving its prognostic accuracy, assess novel CPET-derived metrics of cardiopulmonary fitness (eg, heart rate recovery), and assess possible inter-relationships of CPET-derived metrics with prognostically important comorbidities. For example, lower haemoglobin concentrations are associated with both increased perioperative risk and reduced peak oxygen consumption.<sup>43,44</sup>

In conclusion, preoperative subjective assessment of functional capacity should not be used in clinical practice because it does not accurately identify patients with poor fitness or those at increased risk for morbidity and mortality after major elective non-cardiac surgery. Clinicians could instead consider more objective measures, such as DASI questionnaires and NT pro-BNP testing for assessing perioperative cardiac risk, and perhaps CPET to predict complications after major elective non-cardiac surgery.

#### Contributors

DNW, RMP, MAS, TEFA, BLC, JTG, KET, MPWG, PSM, and BHC contributed to the conception and design of the study. DNW, RMP, MAS, TEFA, ET, AA, BLC, JTG, KET, MPWG, CF, PSM, and BHC contributed to the acquisition, analysis, and interpretation of the data. DNW wrote the first draft of the manuscript. DNW, RMP, MAS, TEFA, ET, AA, BLC, JTG, KET, MPWG, CF, PSM, and BHC revised the manuscript critically for important intellectual content. All authors read and approved the final version of the manuscript.

#### Declaration of interests

TEFA reports grants from the UK Medical Research Council and other funding from the UK National Institute of Health Research (NIHR) Clinical Research Network, and is a committee member of the Perioperative Exercise Testing and Training Society. DNW reports grants and personal fees from Canadian Institutes of Health Research and grants from Heart and Stroke Foundation of Ontario, Ontario Ministry of Health and Longterm Care, and the Ontario Ministry of Research, Innovation and Science. PSM reports grants from the Australian and New Zealand College of Anaesthetists and Monash University. JTG has received research funding from Bayer, Pfizer, and Actelion, and has served on steering, safety, and adjudication committees for Actelion, United Therapeutics, and Bellepheron. RMP reports grants and personal fees from Edwards Lifesciences, Nestle Health Sciences, and BBraun, grants from Intersurgical, and personal fees from Medtronic and GlaxoSmithKline. MPWG reports grants from Sphere Medical and Pharmacosmos, and personal fees from Sphere Medical and Edwards Lifesciences. KET, MAS, ET, BHC, AA, BLC, and CF declare no competing interests.

#### Acknowledgments

This study was supported by grants from the Canadian Institutes of Health Research, Heart and Stroke Foundation of Canada, Ontario



Ministry of Health and Long-Term Care, Ontario Ministry of Research, Innovation and Science, UK National Institute of Academic Anaesthesia, UK Clinical Research Collaboration, Australian and New Zealand College of Anaesthetists, and Monash University (Melbourne, VIC, Australia). DNW is supported by a New Investigator Award from the Canadian Institutes of Health Research. DNW and BHC are partly supported by Merit Awards from the Department of Anesthesia at the University of Toronto. RMP is a Career Development Fellow for the British Journal of Anaesthesia and Royal College of Anaesthetists, and a professor for the UK National Institute for Health Research. TEFA is a clinical research training fellow for the UK Medical Research Council and British Journal of Anaesthesia. MPWG holds the British Oxygen Company Chair of Anaesthesia of the Royal College of Anaesthetists, which is awarded by the UK National Institute of Academic Anaesthesia. We thank the Li Ka Shing Knowledge Institute of St Michael's Hospital (Toronto, ON, Canada) for generously supporting the costs of international trial insurance for this study, and all the participating patients and staff across the 25 study sites.

#### References

- Fleisher LA, Fleischmann KE, Auerbach AD, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014; **130**: e278–333.
- Kristensen SD, Knuuti J, Saraste A, et al. 2014 ESC/ESA Guidelines on non-cardiac surgery: cardiovascular assessment and management: the Joint Task Force on non-cardiac surgery: cardiovascular assessment and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *Eur Heart J* 2014; **35**: 2383–431.
- Melon CC, Eshtiaghi P, Luksun WJ, Wijeyesundera DN. Validated questionnaire vs physicians' judgment to estimate preoperative exercise capacity. *JAMA Intern Med* 2014; **174**: 1507–08.
- Reilly DF, McNeely MJ, Doerner D, et al. Self-reported exercise tolerance and the risk of serious perioperative complications. *Arch Intern Med* 1999; **159**: 2185–92.
- Wiklund RA, Stein HD, Rosenbaum SH. Activities of daily living and cardiovascular complications following elective, noncardiac surgery. *Yale J Biol Med* 2001; **74**: 75–87.
- Hlatky MA, Boineau RE, Higginbotham MB, et al. A brief self-administered questionnaire to determine functional capacity (the Duke Activity Status Index). *Am J Cardiol* 1989; **64**: 651–54.
- Struthers A, Lang C. The potential to improve primary prevention in the future by using BNP/N-BNP as an indicator of silent 'pandcardiac' target organ damage: BNP/N-BNP could become for the heart what microalbuminuria is for the kidney. *Eur Heart J* 2007; **28**: 1678–82.
- Wijeyesundera DN, Pearse RM, Shulman MA, et al. Measurement of Exercise Tolerance before Surgery (METS) study: a protocol for an international multicentre prospective cohort study of cardiopulmonary exercise testing prior to major non-cardiac surgery. *BMJ Open* 2016; **6**: e010359.
- Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004; **240**: 205–13.
- International Surgical Outcomes Study group. Global patient outcomes after elective surgery: prospective cohort study in 27 low-, middle- and high-income countries. *Br J Anaesth* 2016; **117**: 601–09.
- Thygesen K, Alpert JS, Jaffe AS, et al. Third universal definition of myocardial infarction. *Circulation* 2012; **126**: 2020–35.
- Moran J, Wilson F, Guinan E, McCormick P, Hussey J, Moriarty J. Role of cardiopulmonary exercise testing as a risk-assessment method in patients undergoing intra-abdominal surgery: a systematic review. *Br J Anaesth* 2016; **116**: 177–91.
- West MA, Asher R, Browning M, et al. Validation of preoperative cardiopulmonary exercise testing-derived variables to predict in-hospital morbidity after major colorectal surgery. *Br J Surg* 2016; **103**: 744–52.
- Lee TH, Marcantonio ER, Mangione CM, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation* 1999; **100**: 1043–49.
- Ford MK, Beattie WS, Wijeyesundera DN. Systematic review: prediction of perioperative cardiac complications and mortality by the Revised Cardiac Risk Index. *Ann Intern Med* 2010; **152**: 26–35.
- Duceppe E, Parlow J, MacDonald P, et al. Canadian Cardiovascular Society guidelines on perioperative cardiac risk assessment and management for patients who undergo noncardiac surgery. *Can J Cardiol* 2017; **33**: 17–32.
- Pencina MJ, D'Agostino RB, Steyerberg EW. Extensions of net reclassification improvement calculations to measure usefulness of new biomarkers. *Stat Med* 2011; **30**: 11–21.
- Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology* 1990; **1**: 43–46.
- Older P, Smith R, Courtney P, Hone R. Preoperative evaluation of cardiac failure and ischemia in elderly patients by cardiopulmonary exercise testing. *Chest* 1993; **104**: 701–04.
- Young EL, Karthikesalingam A, Huddart S, et al. A systematic review of the role of cardiopulmonary exercise testing in vascular surgery. *Eur J Vasc Endovasc Surg* 2012; **44**: 64–71.
- Carliner NH, Fisher ML, Plotnick GD, et al. Routine preoperative exercise testing in patients undergoing major noncardiac surgery. *Am J Cardiol* 1985; **56**: 51–8.
- McPhail N, Calvin JE, Shariatmadar A, Barber GG, Scobie TK. The use of preoperative exercise testing to predict cardiac complications after arterial reconstruction. *J Vasc Surg* 1988; **7**: 60–68.
- Forshaw MJ, Strauss DC, Davies AR, et al. Is cardiopulmonary exercise testing a useful test before esophagectomy? *Ann Thorac Surg* 2008; **85**: 294–99.
- James S, Jhanji S, Smith A, O'Brien G, Fitzgibbon M, Pearse RM. Comparison of the prognostic accuracy of scoring systems, cardiopulmonary exercise testing, and plasma biomarkers: a single-centre observational pilot study. *Br J Anaesth* 2014; **112**: 491–97.
- Whittle J, Nelson A, Otto JM, et al. Sympathetic autonomic dysfunction and impaired cardiovascular performance in higher risk surgical patients: implications for perioperative sympatholysis. *Open Heart* 2015; **2**: e000268.
- Struthers R, Erasmus P, Holmes K, Warman P, Collingwood A, Sneyd JR. Assessing fitness for surgery: a comparison of questionnaire, incremental shuttle walk, and cardiopulmonary exercise testing in general surgical patients. *Br J Anaesth* 2008; **101**: 774–80.
- Tang WH, Topol EJ, Fan Y, et al. Prognostic value of estimated functional capacity incremental to cardiac biomarkers in stable cardiac patients. *J Am Heart Assoc* 2014; **3**: e000960.
- Coutinho-Myrrha MA, Dias RC, Fernandes AA, et al. Duke Activity Status Index for cardiovascular diseases: validation of the Portuguese translation. *Arq Bras Cardiol* 2014; **102**: 383–90.
- Mclsaac DI, Bryson GL, van Walraven C. Association of frailty and 1-year postoperative mortality following major elective noncardiac surgery: a population-based cohort study. *JAMA Surg* 2016; **151**: 538–45.
- Rodseth RN, Biccard BM, Le Manach Y, et al. The prognostic value of pre-operative and post-operative B-type natriuretic peptides in patients undergoing noncardiac surgery. B-type natriuretic peptide and N-terminal fragment of pro-B-type natriuretic peptide: a systematic review and individual patient data meta-analysis. *J Am Coll Cardiol* 2014; **63**: 170–80.
- Botto F, Alonso-Coello P, Chan MT, et al. Myocardial injury after noncardiac surgery: a large, international, prospective cohort study establishing diagnostic criteria, characteristics, predictors, and 30-day outcomes. *Anesthesiology* 2014; **120**: 564–78.
- Devereaux PJ, Xavier D, Pogue J, et al. Characteristics and short-term prognosis of perioperative myocardial infarction in patients undergoing noncardiac surgery: a cohort study. *Ann Intern Med* 2011; **154**: 523–28.
- Beattie WS, Wijeyesundera DN, Chan MTV, et al. Survival after isolated post-operative troponin elevation. *J Am Coll Cardiol* 2017; **70**: 907–08.
- Moonesinghe SR, Harris S, Mythen MG, et al. Survival after postoperative morbidity: a longitudinal observational cohort study. *Br J Anaesth* 2014; **113**: 977–84.
- Devereaux PJ, Mrkobrada M, Sessler DI, et al. Aspirin in patients undergoing noncardiac surgery. *N Engl J Med* 2014; **370**: 1494–503.
- Myles PS, Smith JA, Forbes A, et al. Stopping vs. continuing aspirin before coronary artery surgery. *N Engl J Med* 2016; **374**: 728–37.

- 37 Sheth T, Chan M, Butler C, et al. Prognostic capabilities of coronary computed tomographic angiography before non-cardiac surgery: prospective cohort study. *BMJ* 2015; **350**: h1907.
- 38 Myles PS, Leslie K, Chan MT, et al. The safety of addition of nitrous oxide to general anaesthesia in at-risk patients having major non-cardiac surgery (ENIGMA-II): a randomised, single-blind trial. *Lancet* 2014; **384**: 1446–54.
- 39 Kopec M, Duma A, Helwani MA, et al. Improving prediction of postoperative myocardial infarction with high-sensitivity cardiac troponin T and NT-proBNP. *Anesth Analg* 2017; **124**: 398–405.
- 40 Ndumele CE, Matsushita K, Sang Y, et al. N-Terminal pro-brain natriuretic peptide and heart failure risk among individuals with and without obesity: the Atherosclerosis Risk in Communities (ARIC) Study. *Circulation* 2016; **133**: 631–38.
- 41 Gregg LP, Adams-Huet B, Li X, et al. Effect modification of chronic kidney disease on the association of circulating and imaging cardiac biomarkers with outcomes. *J Am Heart Assoc* 2017; **6**: e005235.
- 42 Tromp J, Khan MA, Klip IT, et al. Biomarker profiles in heart failure patients with preserved and reduced ejection fraction. *J Am Heart Assoc* 2017; **6**: e003989.
- 43 Fowler AJ, Ahmad T, Phull MK, Allard S, Gillies MA, Pearse RM. Meta-analysis of the association between preoperative anaemia and mortality after surgery. *Br J Surg* 2015; **102**: 1314–24.
- 44 Otto JM, O'Doherty AF, Hennis PJ, et al. Association between preoperative haemoglobin concentration and cardiopulmonary exercise variables: a multicentre study. *Perioper Med (Lond)* 2013; **2**: 18.